GROWING CARNIVOROUS PLANTS IN THE UK

MARCELLO CATALANO • via Ronchi 2 • 20134 • Milano • Italy • rafflesiana@yahoo.com

Keywords: cultivation: Great Britain — live collections: Kew.

Here is the story of how I brought some Mediterranean atmosphere to the green climate of England.

A little while ago, I decided to travel and work abroad, so I spent April through June 2001 working at Kew Gardens, in London. It was a great opportunity for me to work in what is maybe the most important botanical garden in the world.

Kew Gardens were founded in 1759. Today 550 people work there, of which 200 are gardeners, 150 are scientific staff, and the others are technical and support staff. The library contains more than 130,000 volumes and 4000 periodicals. The herbarium, one of the largest in the world, contains seven million specimens. The plant and seed collections are the largest in the world.

The Garden has about 40,000 kinds of plants on view for the public—the rest are grown in the more secure environment of the Garden's nurseries. There are three greenhouses open to the public: the temperate house, the tropical house, and the Conservatory. The Conservatory is for the most precious and interesting collections (cacti, carnivorous plants, orchids, ferns, bromeliads, aroids, etc.).

Not accessible to the public is a temperate nursery and a tropical nursery. Each nursery is about as large as the three greenhouses all together.

When I arrived at Kew in April, the specimens on display were very large and nice (such as their *Sarracenia flava* var. *rugelii*), but apart from a large *Utricularia reniformis*, the most precious stuff was in the nursery, far from danger by being damaged by too curious people or kid's hands. In the previous year almost all of the carnivorous plant collection had been lost because of the poor quality of water. However, Kath King, from Perth Australia, was now in charge of the carnivorous plants and was working to rebuild the carnivorous plant collection.

The tropical nursery contains a coolhouse for Sarracenia, Drosera, Pinguicula, Cephalotus, Dionaea and Drosophyllum, and a warm house for Nepenthes, Heliamphora and some tropical Utricularia. Darlingtonia and temperate Pinguicula (such as P. grandiflora) are grown outside. Byblis gigantea, P. primuliflora and some other Nepenthes and Heliamphora are kept in a well-lighted bromeliad greenhouse.

In the warm house, *Nepenthes* cover a full bench almost 2 meters large and about 10 meters long. I remember best the specimens of *N. hamata* (pitchers 15 cm long), *N. pervillei* (plant is 30 cm large), *N. macfarlanei* (pitchers 25 cm long), *N. rajah* (plants 15 cm large), *N. northiana*, *N. merrilliana* and *N. madagascariensis*. There are many other interesting species including *N. hirsuta*, *N. treubiana*, *N. veitchii*, *N. reinwardtiana* (80 cm tall), *N. bongso*, *N. ramispina*, *N. sanguinea*, *N. maxima*, *N. ventricosa*, *N. tobaica*, *N. mirabilis* (1-2 meter plants), *N. thorelii*, *N. × mixta* (1-2 meters tall), and *N. tobaica* (80 cm tall), etc. Before you have a heart attack, you should know there are no giants. Still, even if they need a lot of love and care to become 2-3 meters climbers, they are healthy, large and strong.

The warmhouse also has *U. alpina*, *U. calycifida* and *U. prehensilis*, *U. reniformis*, *U. praelonga*, *U. tricolor*, *U. pubescens* and *U. bisquamata*.

The coolhouse has 10-20 specimens of every Sarracenia species, about fifty

Dionaea, about thirty Drosera species (tuberous, pygmies and others; 1-10 specimens of each). There is also U. sandersonii, U. dichotoma, U. livida, and U. bisquamata from Betty's Bay; other plants include Mexican Pinguicula (P. laueana, P. gypsicola, P. esseriana), P. primuliflora, P. lusitanica, a flowering Cephalotus (20 cm across), 3-4 Drosophyllum, Byblis gigantea and an elegant, 25 cm tall Roridula dentata.

The best, red *Heliamphora nutans* and *H. heterodoxa* are in the *Nepenthes* house. More *H. nutans*, *H. minor*, a nice flowering *B. gigantea*, *N. rajah* and *N. macfarlanei* are quite happy in the bromeliad house, which is a well-lit, relatively low humidity environment.

On 21 April Kath King and I attended the annual general meeting of the UK Carnivorous Plant Society, held at the Reading University, near London.

I met a lot of people from the Carnivorous Plant Society, and saw the University collection, wealthy with some huge *S. minor*, a lot of healthy *Drosera*, trays filled with red *Dionaea*, a deep purple *Drosera binata* and other colorful plants. After some lovely tea and biscuits we listened to some lectures; I read a short speech encouraging collaboration of carnivorous plant societies. For example, I encouraged collaboration between the UK Carnivorous Plant Society and Kew Gardens. At the end, Kath King agreed in promoting a better and more intense collaboration between the Kew Gardens and the Carnivorous Plant Society, and in August there will be an open day in the Kew tropical nursery specifically for the Carnivorous Plant Society.

On the 25 of May I visited the Chelsea Flower Show (see Figures 1, 2). (My ticket was provided free, compliments of the Carnivorous Plant Society.) I spent most of the day talking with the growers in the three carnivorous plant stands: the UK Carnivorous Plant Society (silver medal winner), South West Carnivorous Plants (silver medal winner) and Hampshire Carnivorous Plants (gold medal winner). You can really feel the 25 years of the UK Carnivorous Plant Society's history, talking with the kind and always available people. They know everything! In few hours I learned how to feed *Utricularia* with nematodes to make them flower (Paul McKeown, seed bank manager), I listened to different opinions about Don Schnell's classification of the different varieties of *S. flava* (Matthew Soper of Hampshire Carnivorous Plants, Chris Crow of *Sarracenia* Nurseries), I know a lot more about neon lamp brands (Alistair Pearce of South West Carnivorous Plants), and heard about the history of the UK Carnivorous Plant Society (Steve Cottel, chairman). They also provided me ideas (Alan Haines, web builder and Malcolm Goddard, newsletter editor) on how to improve the Italian Carnivorous Plant Society.

The specimens on display were fabulous: *U. reniformis* and *U. longifolia* var. forgetiana covered with flowers, perfect and giant *S. psittacina*, *S. flava*, *S. minor*, shining, emerald-like Darlingtonia, *P. moranensis* with white flowers, *P. crystallina*, *P. laueana* and many others with plenty of flowers. Also present were vast numbers of tuberous and pygmy Drosera (including a magnificent *D. gigantea*). While Cephalotus, Heliamphora and Nepenthes are my main interest—and the specimens of them there were beautiful—I almost ignored them, as I was completely taken by the beauty of all the rest.

I helped dismantle the stand and load the plants onto a truck at the end of the show. It was half past nine in the evening when, tired by the intense day but enthusiastic for all the growing information I learned, I returned home.

While I worked at Kew, Phil Wilson and Paul McKeown often visited Kew to help Kath King with the identification and the cultivation of the plants. As a bonus, they often brought spare plants. I was present during one of these visits. I remained in silence while they found tens of incorrectly labelled *Drosera* pots. All those sun-



Figure 1: The Carnivorous Plant Society's display at Chelsea flower show, 2001. Photo by Alan Haines.



Figure 2: The Carnivorous Plant Society's display at Chelsea flower show, 2001. Photo by Alan Haines.

dews that to me were simply all *D. spathulata* (sundews are not my specialty), were recognized with amazing ability and quickness. We spoke about Kew's *Sarracenia rubra* subspecies, about a strange *N. macfarlanei* that may actually be a hybrid, about *Pinguicula ehlersiae* and how often it is sold as *P. cyclosecta*. Over biscuits and a cup of tea, they related what Kew's carnivorous plant collection was like in the past, they recounted tales of *Drosera* floating on the water in some bogs on the Tepuis, they described floating *Pinguicula primuliflora* in some American lakes, and discussed horticulturists performing such feats as growing huge *H. tatei* grown as an epiphyte on a slab of bark and some lucky people who are growing rope-size *Aldrovanda* in ponds where the conditions are perfect.

On 14 June I met the famous Martin Cheek in the Kew Herbarium. Martin is the professor who has done a great deal of work in the study of Nepenthes taxonomy. If you look carefully in your carnivorous plant books, after names such as N. argentii, N. aristolochioides, N. danseri, N. diatas, N. lamii and N. macrophylla you will read "Jebb and Cheek". He is that Cheek. Martin has been working at the Kew Herbarium on a long-time project classifying South African flora, but my visit gave him an opportunity to return to one of his favourite genera: Nepenthes. Martin is a very kind person, he always speaks slowly with a low voice—almost whispering and hypnotizing. When he speaks about Nepenthes you can see in his eyes the shine of someone who has, for his passion, the curiosity and the enthusiasm of a child. He delighted me for an hour of his precious time with some words about the differences between N. burkei and N. ventricosa, N. stenophylla and N. faizaliana, and N. thorelii and N. smilesii (even showing me dried specimens), and described the habitat of the two amazing species N. merrilliana and N. northiana. He showed me material that Peter Taylor used to write his monograph on Utricularia (including even his microscope!).

The 22 and 23 of June were unforgettable. I took the bus from London and after 3 and a half hours I arrived at Culmstock, Devon, in the evening. I was to visit







Figure 3: Mike King's Sarracenia flava var. rubricorpora, in his 10 meter long greenhouse. Photo by Mike King.

Alistair and his wife Jenny Pearce and their nursery "South West Carnivorous Plants". Al and Jenny are two delightful people—I brought them an Italian wine as a gift, and they offered me a dinner in a local pub. Before it grew too dark, I saw his vast and diverse collection, and was particularly impressed by hundreds of colorful Sarracenia, a large tray of U. sandersonii, a huge N. veitchii (the highland form, with striped peristome), and some giant Darlingtonia with their bright heads shining at the sunset light. We spent the whole evening speaking about carnivorous plants, sharing photos, and surfing the web. Since I was visiting with Phil Wilson the next day—only half an hour from Culmstock—Alistair and Jenny kindly offered for me to sleep there, and then to accompany me to Phil's house the next day.

Finally, it was time to sleep. As I dropped into sleep I felt all the romantic beauty of traveling; while the moonlight came through the curtains and a dog was barking somewhere, I realized that I was sleeping on a divan at Culmstock, a small village in the southwest of England.

The following morning we arrived at Phil Wilson's home in Somerset (where also lives Adrian Slack). I gave Phil the second bottle of Italian Moscato and I took some pictures of his extensive collection. There are a lot of rare specimens of many different carnivorous plants: Roridula gorgonias, U. humboldtii, U. multifida in flower, a lidless clone of S. flava, S. leucophylla 'Schnell's Ghost', etc. I saw some people I had met earlier at the Chelsea Flower Show, and I helped them to put new UK CPS bulletins in envelopes for mailing. Paul McKeown gave me a lift back to London (on the way back I saw Stonehenge!). This drive with Paul was possibly the most instructive conversation of the whole trip as for hours we discussed many different carnivorous plant growing techniques.

It was 7 July when my tired limbs could finally relax—the Kew internship was over and I had more free time to enjoy England. Alan Haines helped me to not get lost with the English trains, and with another bottle of Moscato in hand, we visited Mike King's collection near Birmingham. It was the most beautiful Sarracenia collection I had ever seen (see Back Cover and Figure 3). For such a large collection there was a perfect order: all the plants were labelled, the greenhouse windows were clean, the concrete floor was free of rubbish, old pots and compost. All the Sarracenia were strong, healthy and colorful, all the dead leaves and spent flowers were trimmed away. Mike did not use greenhouse fans or shades. Instead, the roof windows automatically opened when the temperature rises too much. The plants were obviously happy about the full sun, plentiful rain water and fresh air. Mike was a perfect host, he ran from one guest to the other, answering questions and showing off this or that plant. The food was good and plentiful, And, after showing me around his home—including his supplies of peat and perlite, and a barrel system he uses to collect rain water—he confided to me that these plants are his life. I thought back to when I had first met Mike during a previous meeting of growers and how he always enjoyed showing off pictures of his wonderful Sarracenia...I think that anyone would do the same if they had such fine specimens!

On 14 July I travelled by train to meet Matthew Soper, owner of "Hampshire Carnivorous Plants" and winner of a gold medal at the Chelsea Flower Show. For Matthew, I brought along a bottle of light sweet wine from Reggio Emilia. His collection was vast—hundreds of Sarracenia species and hybrids and a few Drosera. But what most attracted me were his Nepenthes and Utricularia. With pop music dancing in the air, Matt showed me with pride his thirteen year-old N. rajah with a beautiful pitcher about 25 cm long! I saw some beautiful N. macrophylla (20 cm pitchers), highland and lowland forms of N. veitchii (20 cm pitchers), N. inermis (10 cm pitchers), N. hirsuta (20 cm pitchers), N. lowii (10 cm pitchers), a wonderful N. veitchii × maxima (25 cm pitchers) and many others. He also showed me his

Heliamphora ionasi (rather small, but you know how slow this species grows), H. tatei (10 cm tall) and an unrecognizable, strange Heliamphora. I asked him if it was the new, recently discovered H. hispida, but he was not sure, as he received it just last year labelled only as a "new species." There were some huge Darlingtonia, among which I saw a clone covered with smaller and more numerous windows than usual. He had attractive clones of Cephalotus that, just like movie stars, looked stressed because of the too many shows they had taken part in. Matt also had a shocking, giant S. flava var. rubricorpora—I never before saw a pitcher as large as that! There was a large tray filled with *Utricularia* species, most of them in flower, among which I saw the precious *U. tricolor*, *U. tridentata*, *U. longifolia*, *U. pubescens*, U. nephrophylla, U. blanchetii and some astonishing U. praelonga full of scapes covered with huge yellow flowers. Over tea, we looked at some very interesting Nepenthes pictures he took in nature, and three hours went by like they were few minutes. As a note of particular interest, we spoke about how Thailand, Cambodia, Vietnam and Indochina are rarely explored, and how there may be plenty of new species just waiting for brave taxonomists.

The very next day, I made my last visit to a UK grower, meeting Paul McKeown in his house near Watford, north of London. (Paul does not like wine, so I brought him some beer.) His Sarracenia collection was suffering because of root mealybugs infesting his plants after the Chelsea Flower Show. (It seemed that all the growers complained about the loss or worsening of many carnivorous plant specimens after the Flower Show.) Paul had great plants, like flowering Genlisea violacea and Genlisea hispidula, and large specimens of U. quelchii, U. endresii, and U. nelumbifolia. His 10 year-old wonderful, huge Darlingtonia was fantastic. But what nearly mesmerized me is his highland Nepenthes collection. He had an impressive N. veitchii vine (1 meter long), an eight year-old N. rajah (along with a few "small" specimens 25 cm in diameter), N. macrophylla (pitcher: 20 cm), N. aristolochioides (pitcher: 8 cm), N. sanguinea (pitcher: 15 cm), N. maxima (pitcher: 30 cm), N. fusca (pitcher: 20 cm), N. bongso (pitcher: 20 cm), N. spectabilis (pitcher: 20 cm), N. mikei (pitcher: 12 cm), N. glabrata (pitcher: 5 cm), and N. sibuyanensis (pitcher: 12 cm). In addition, he had many smaller but healthy specimens of N. pilosa, N. burbidgeae, N. lowii, N. hamata, N. ovata and others. Paul told me more than once that the problems with highland Nepenthes were often due to inadequately-developed root systems in plants coming from micropropagation. Another important point was that (apart the usual exceptions like N. alata, N. ventricosa, N. khasiana, and N. maxima), these species really did poorly in overly warm conditions. (Paul is using 50% shade and a humidifier that sprays cold fog about every 20 minutes. The temperature, even in summer, rarely goes over the 25°C (77°F) and the night minimum is 12°C (54°F).) Paul said that these two points were more important than what kind of soil is used, or how much humidity and light the plants were given. Once the small plants have developed a strong root system in the cool environment, they will grow quickly and bigger, just like lowland species. The evidence was before me! This was the first time I saw all these highland species growing so big; even the small plants were healthy, with every leaf and pitcher bigger than the previous ones.

Well, that concludes my experiences in the UK. I saw wonderful plants, I learned a lot of new growing tips and I met many friendly, interesting and experienced growers that I wish to thank for their kindness. I hope they enjoyed our discussions as much as I did, and I hope one day, if anyone of them ever come to Italy, we will be able to show off as many beautiful and rare specimens as I saw on my UK travels.

CARNIVOROUS PLANT



Journal of the International Carnivorous Plant Society www.carnivorousplants.org

Volume 31, Number 4 December 2002



Front Cover: Drosera cuneifolia flowers. See article on page 100.

Back Cover: Mike King's giant *Sarracenia flava* var. *rugelii*. Photo by Mike King. See article on page 120.

Carnivorous Plant Newsletter is dedicated to spreading knowledge and news related to carnivorous plants. Reader contributions are essential for this mission to be successful. Do not hesitate to contact the editors with information about your plants, conservation projects, field trips, or noteworthy events. Contributors should review the "Instructions to Authors" printed in the March issue of each year. Advertisers should contact the editors. Views expressed in this publication are those of the authors, not the editorial staff.

All correspondence regarding dues, address changes and missing issues should be sent to the Membership Coordinator at the ICPS. Do not send such correspondence to the editors. Checks for subscriptions and back issues should be made to the ICPS in US funds. Dues for 2002 are \$25.

ICPS, Inc. PMB 330

3310 East Yorba Linda Blvd. Fullerton, CA 92831-1709, USA icps@carnivorousplants.org

President David Gray, email: david@carnivorousplants.org
Vice President Carl Mazur, email: carl@carnivorousplants.org
Secretary Cindy Slezak, email: cindy@carnivorousplants.org

Seed Bank John Brittnacher, email: john@carnivorousplants.org, seedbank listed in this issue.

Editors:

Barry A. Rice, P.O. Box 72741, Davis, CA 95617, USA, email: barry@carnivorousplants.org Jan Schlauer, Zwischenstr. 11, D-60594 Frankfurt, Germany, email: jan@carnivorousplants.org Page Layout: Steve Baker, email: steve@carnivorousplants.org

Date of effective publication of the September 2002 issue of Carnivorous Plant Newsletter: 6 September 2002.

The ICPS is the International Cultivar Registration Authority (ICRA) for cultivated carnivorous plants according to The International Code For The Nomenclature of Cultivated Plants. Send relevant correspondence to the ICPS, Inc.

PUBLISHER: ICPS, Inc., Fullerton, California. Published quarterly with one volume annually. Desktop Publishing: Steve Baker, 5612 Creek Point Drive, Hickory, NC 28601. Printer: Kandid Litho. Logo and masthead art: Paul Milauskas. Dues: \$25.00 annually. Reprints available by volume only © 2002 Carnivorous Plant Newsletter. All rights reserved. ISSN #0190-9215.

